

Endocrine Pharmacology

GALGOTIAS
UNIVERSITY

Disclaimer

All the content material provided here is only for teaching purpose.

GALGOTIAS
UNIVERSITY


Endocrine Pharmacology

Hormone is a substance of intense biological activity that is produced by specific cells in the body and is transported through circulation to act on its target cells.

These substances are secreted by one cell (Secretary cell) and perform action on other cells (Target Cell).

The word endocrine means '**Ductless gland**' i.e.- this secretion directly goes to the blood from the secretary cell.


Endocrine glands:



Hypothalamic & Pituitary Hormones

The output of the hypothalamus-pituitary unit regulates the function of the thyroid, adrenal and reproductive glands and also controls somatic growth, lactation, milk secretion and water metabolism.

Hypothalamic Hs can have effect of stimulating or inhibiting the release of ant. Pit. Hs. Called **RELEASING HORMONES “RH”** or **INHIBITING HORMONES “IH”** respectively, reflecting their influence on ant. Pit. Hs.



The Pituitary Gland is divided into 2 areas, *with separate types* of hormone production.

Anterior pituitary

- Growth Hormone (**GH**)
- Thyroid-stimulating Hormone (**TSH**)
- Adrenocorticotropin (**ACTH**)
- Follicle-stimulating Hormone (**FSH**),
- Leutinizing Hormone (**LH**),
- Prolactin

Posterior pituitary


- Oxytocin and
- Antidiuretic hormone (**ADH**)

Control of Pituitary glands by Hypothalamus

1 Releasing and inhibiting hormones are released from hypothalamic neurons into the hypothalamo-pituitary portal system.

2 Hypothalamic-releasing and hypothalamic-inhibiting hormones are carried down the pituitary stalk by the hypothalamopituitary portal system.

3 Hypothalamic-releasing and hypothalamic-inhibiting hormones increase or decrease, respectively, the release of anterior pituitary hormones into general circulation.



1 Oxytocin and vasopressin are synthesized in the paraventricular and supraoptic nuclei of the hypothalamus.

2 Oxytocin and vasopressin are carried by axonal transport down the pituitary stalk.

3 Oxytocin and vasopressin are released into general circulation from terminal buttons in the posterior pituitary.

Anterior Pituitary Hormone:

Growth Hormone

Derived from the somatotroph cells

Its secretion is controlled by GHRH and somatostatin

- GH secretion is high in newborn, decreasing at 4 yr to an intermediate level, which is then maintained until puberty, when there is further decline
- Insulin-like growth factor 1 (IGF-1) released from the liver inhibits GH secretion by stimulating somatostatin secretion from the hypothalamus

Growth Hormone Activity

- Increases plasma free fatty acids (source of energy for muscle tissue)
- Increases hepatic glucose output
- Decreases insulin sensitivity in muscle
- Is protein anabolic hormone

Growth Hormone Deficiency

- In childhood: short stature and adiposity, hypoglycemia.
- Adults : generalized obesity, reduced muscle mass.

Growth Hormone Excess

- In adults causes acromegaly
- If this occurred before the long bone epiphyses close, it leads to the rare condition, gigantism

Thyroid-stimulating Hormone (TSH)

- Also called thyrotrophin
- Stimulates secretion of thyroid hormone & growth of thyroid gland.

Adrenocorticotropin (ACTH)

- Stimulates cortisol secretion by the adrenal cortex & promotes growth of adrenal cortex

Follicle –stimulating hormone (FSH)

- Females: stimulates growth & development of ovarian follicles, promotes secretion of estrogen by ovaries.
- Males: required for sperm production

Leutinizing hormone (LH)

- Females: responsible for ovulation, formation of corpus luteum in the ovary, and regulation of ovarian secretion of female sex hormones.
- Males: stimulates cell in the testes to secrete testosterone

Prolactin

- Secreted by lactotroph cells of the ant. Pit., which increase in number during pregnancy.
- Its secretion is stimulated by estrogen
- Females: stimulates breast development and milk production.
- Males: involved in testicular function

Posterior pituitary Hormones

Oxytocin


- It is synthesized in the hypothalamus & transported to the post. Pit.
- It is an effective stimulant of uterine contractions & is used intravenously to induce or reinforce labor .
- Induces the release of milk
- Suckling sends a message to the hypothalamus via the nervous system to release oxytocin, which further stimulates the milk glands

Vasopressin (antidiuretic hormone ADH)

- The function of ADH is to increase water conservation by the kidney.
- ADH release increases if blood pressure falls or blood becomes too salty.
- ADH causes peripheral blood vessel constriction to help elevate blood pressure .

Regulation of Hormonal secretion by Hypothalamus and Pituitary Gland

Endocrine Glands



Thyroid Hormones

Triiodothyroxine T3

Tetraiodothyroxine T4

Calcitonin(Parathyroid)


A large, faint watermark logo of Galgotias University is centered on the page. It features a stylized 'G' composed of three curved, overlapping bands in shades of yellow, blue, and red, all contained within a larger, light-colored circular shape.

**GALGOTIAS
UNIVERSITY**


Synthesis of T3 and T4

It involves the following steps:

- Iodide Uptake
- Oxidation and Iodination
- Coupling
- Storage and release
- Peripheral conversion of T4 to T3



Regulation of T3 and T4



Physiological action

Metabolism

- Glucogenolysis and gluconeogenesis → increase blood glucose level. ↑ lipolysis.
- Increase BMR, induction of haemopoiesis and increase peristalsis of GIT.

Hypothyroidism: Deficiency of thyroid hormones revealed the following effects:

In the Children: *Cretinism* (mentally retarded, dwarf).

In the adult: Myxoedema: myxomatous tissues get deposited on the face and lead to weight gain, decrease BMR, heart rate, body temperature. and in the females, additionally cause amenorrhoea.

Endemic goiter: Increase the size of thyroid gland.

Antithyroid Drugs

- A. Inhibit hormone synthesis:** Propyluracil, Methimazole, Carbimazole
- B. Inhibit iodide trapping (ionic inhibitors):** Thiocynates ($-\text{SCN}$), Perchlorate (ClO_4^-), Nitrates (NO_3^-)
- C. Inhibit hormone release:** Iodine, Iodide of Na and K, organic iodides
- D. Destroyed thyroid tissue:** Radioactive iodine (^{131}I , ^{125}I , ^{123}I)

Thioamides

MOA: Bind to the thyroid peroxidase and prevent oxidation of iodide/iodotyrosyl residues, thereby;

- (i) Inhibit iodination of tyrosine residues in thyroglobulin
- (ii) Inhibit coupling of iodotyrosine residues to form T3 and T4.

Pharmacokinetics:

- All antithyroid drugs are quickly absorbed orally
- widely distributed in the body, enter milk and cross placenta
- Metabolized in liver
- Excreted in urine primarily as metabolites

Adverse effects:

- Hypothyroidism and goiter

Parathyroid Hormones:

These hormones are secreted by Parathyroid gland situated just behind the Thyroid gland.

- Parathyroid Hormones is secreted by the chief cells of the parathyroid glands as polypeptide containing 84 amino acids .
- It acts to regulate the concentration of calcium (Ca^{2+}) and Phosphate in the blood.
- Important in bone remodeling, which is an ongoing process in which bone tissue is alternately resorbed & rebuilt over time.

Function of Parathyroid Hormones (Pth)

- Increase the osteoclast cells activity → resorption of bone Ca^{2+} → increase blood Ca^{2+} level → hypercalcemia.
- Increase the intestinal absorption of Ca^{2+} → increase blood Ca^{2+} level
- Increase Po_4^{3-} excretion.

Function of calcitonin: Opposite to Parathyroid hormone.

Hypoparathyroidism Low plasma level of Ca^{2+} and tetany, laryngospasm.

Hyperparathyroidism Hypercalcemia, decalcification of bones (osteoporosis) osteitis fibrosa cystic and renal stone.

Vitamin D

- **D2** Calciferol (ergocalciferol): Obtained from plant sources–food.
- **D3** Cholecalciferol: Synthesized in the skin under the influence of UV radiation.
- **D3** → Calcitriol (active form of D3 –1, 25 dihydroxy cholecalciferol) which is responsible for physiological function and which is similar to parathyroid hormone.
- **Uses of vitamin D:** In rickets, senile or postmenopausal osteoporosis, hypoparathyroidism, and Fanconi syndrome.

References

1. **Tripathi KD. 'Essentials of Medical Pharmacology', 6th edition, Jaypee Brothers Medical publications (P) Ltd., New Delhi, 2003.**
2. **Mohan H. 'Text book of Pathology', 4th edition, Jaypee Brothers Medical publications (P) Ltd., New Delhi, 2004.**
3. **Dale, M M, H P. Rang, and Maureen M. Dale. '*Rang & Dale's Pharmacology*', 7th edition. Edinburgh: Churchill Livingstone, 2007.**
4. **Whalen, Karen, Richard Finkel, and Thomas A. Panavelil. *Lippincott Illustrated Reviews: Pharmacology*. 6th ed. Philadelphia, PA: Wolters Kluwer, 2015.**
5. **Satoskar RS, Ainapure SS, Bhandarkar SD, Kale AK, 'Pharmacology and pharmacotherapeutics', 14th edition, Popular Prakashan, Mumbai, 1995.**